



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MAY 15.

REV. H. C. MCCOOK, D. D., Vice-President, in the chair.

Eighteen persons present.

The deaths were announced of Caleb Cope, a member, on the 12th, inst. and Dr. Gerhard Vom Rath, a correspondent, April 23.

A paper entitled "Notes on new species of Orb-weaving Spiders." By Rev. Henry C. McCook was presented for publication.

Notes on the Relations of Structure and Function to Color Changes in Spiders.—Rev. Dr. HENRY C. MCCOOK submitted the following remarks on color changes in spiders, which he wished to be understood as in part, at least, tentative. They were intended to evoke suggestions and helpful information from members of the Academy and others, rather than to present final conclusions on a most interesting subject.

I. On the Relation of Structure to Color he observed that:—

1 The color of young spiders is almost without exception light yellow, or green, whitish or livid, tints that blend very well with the prevailing greens of foliage, young twigs and the grays of bark of trees, of rocks and soil. This is due largely to the fact that the tissues are at that time translucent, allowing a free play of light through them. The effect is also, probably, caused by the absence of food in the alimentary tract and lack of distribution of nutriment throughout the system.

As young spiders advance in age the color deepens, which is caused no doubt by gradual hardening of the tissues, thus making them more opaque. Up to this period no food has been taken, hence the absence of food alone is not sufficient to account for the light colors of the first stages after exode. Yellows and browns in various tints occur at this period, and in some cases, not generally he believed, color patterns which are characteristic of the various species in adult life begin to appear with more or less distinctness, or at least suggestively. It is not until sedentary spiderlings have established themselves upon their own webs, and so to speak, have set up housekeeping for themselves, that the characteristic colors of the species begin to appear with any positive degree of distinctness.

2 As the spiders further advance in age and make their successive moults, various color changes may be noted. Immediately after moulting the color is always lighter, which is probably due to the fact that the harder skin, just cast off, prevented the passage of light through the tissues. The new skin is probably thinner, and more translucent. Dr. McCook believed that moulting produces changes in color patterns of a very decided kind, at least in certain species. Apparently some organic change occurs which is the cause of this phenomenon.